

Patients currently receiving Strontium

Has been on therapy for less than 3 years.

Has been on therapy for greater than 3 years.

No past history of bisphosphonate

One bisphosphonate tried in the past for duration of less than 5 years

More than one bisphosphonate in past history

Has patient sustained a low trauma fracture whilst on strontium

Initiate Alendronate or Risedronate

Offer a second bisphosphonate
OR
Refer to FLS/ Barts Health Osteoporosis Service

Refer to FLS/ Barts Health Osteoporosis Service

No

Yes

Review after 5 years

Stop strontium and continue calcium and vitamin D

Refer to FLS/ Barts Health Osteoporosis Service

DEXA scan after 12-18 months

If BMD drops manage according to primary care osteoporosis pathway

Refer and contact details to Fracture Liaison Service (FLS)

Osteoporosis / Fracture Prevention Nurse
Senior Sister Dee Tomasso

Tel No: 01708 435000 ext 3219
Email: Dee.tomasso@bhrhospitals.nhs.uk



Osteoporosis/Fracture Prevention Service
Referral

Date of referral:

Patient Name:

Hospital No:

DOB:

Tel No:

Fracture – Wrist† Humerus† Vertebrae† Other†

Reason for referral

Referred by (Please Print Name)_____

Signature_____

Send to:

Osteoporosis / Fracture Prevention Nurse

Senior Sister Dee Tomasso

Team 1 Room 28 Out Patients Department,

Queens Hospital,

Rom Valley Way,

Romford,

Essex,

RM7 0AG

Contact Details:

Tel No; 01708 435000 ext 3219



Bart's Health Osteoporosis Service
Referral

Date of referral:

Patient Name:

Hospital No:

DOB:

Tel No:

Reason for referral

Referred by (Please Print Name)_____

Signature_____

Send to:
Osteoporosis Service
FAO Dr Simon Donnelly
The Silverthorn Centre,
Chingford,
London,
E4 6UN

